## One single procedure for your crown cementations

Zirconia, Lithium Disilicate or Metal. No primers required. One single procedure regardless of the restorative material.



Conditioning the crown.\* \* Refer to "Conditioning the prosthetic



Dispense an equal amount of Paste A & B. Mix Paste A & B for 10 seconds.



Apply the cement to the crown.



Place the crown.




Light-cure for 2 to 5 seconds or chemical-cure for 2 to 4 minutes, then remove the excess cement.

Maintain isolation for 5 minutes.



Follow the instructions for use of the restoration material. In the absence of specific instructions, we recommend the following procedure:













Blast with alumina powder (30~50 µm), then ultrasonic clean and dry.



PANAVIA™ SA Cement Universal **Automix** 

- · Universal (A2)
- •Translucent Content: 1 syringe
- Paste A and B: 8.2 g (4.6 mL), Mixing Tip: 20 tips



PANAVIA™ SA Cement Universal Handmix

- · Universal (A2)


- Translucent
  Content: 1 syringe:
  Paste A and B: 9.2 g (5 mL)



Website www.kuraraynoritake.com

US Distributed by Kuraray America, Inc. 33 Maiden Lane, 6th Floor New York, NY 10038

Kuraray Europe, GmbH Philipp-Reis-Str. 4 65795 Hattersheim am Main, Germany



- · Printed color can be slightly different from actual color.

"PANAVIA", "AVENCIA" and "CLEARFIL" are trademarks of Kuraray Co., Ltd. "KATANA" is a trademark of NORITAKE CO., LIMITED.











## **PANAVIA SA Cement Universal**

## SIMPLE, CLEAN AND VERSATILE

FORGET ABOUT PRIMERS



Kuraray Noritake Dental Inc.